



Detection of *Salmonella* from animal sources in South Africa between 2007 and 2014

**Authors:**

Awoke K. Gelaw¹ 
 Palesa Nthaba¹
 Itumeleng Matle¹ 

Affiliations:

¹Agricultural Research Council, Irene, South Africa

Corresponding author:

Awoke Gelaw,
 kidanemariam@arc.agric.za

Dates:

Received: 27 Nov. 2017

Accepted: 16 July 2018

Published: 07 Nov. 2018

How to cite this article:

Gelaw, A.K., Nthaba, P. & Matle, I., 2018, 'Detection of *Salmonella* from animal sources in South Africa between 2007 and 2014', *Journal of the South African Veterinary Association* 89(0), a1643. <https://doi.org/10.4102/jsava.v89i0.1643>

Copyright:

© 2018. The Authors.
 Licensee: AOSIS. This work is licensed under the Creative Commons Attribution License.

Retrospective laboratory-based surveillance was conducted on *Salmonella* serotypes isolated from various animal species from 2007 to 2014 at the Agricultural Research Council, Onderstepoort Veterinary Research Institute, South Africa. During the surveillance period, 1229 salmonellae isolations were recorded. Around 108 different serotypes were recovered from nine different food and non-food animal host species. The three most common serotypes were *Salmonella enterica* subspecies *enterica* serotype Heidelberg ($n = 200$), *Salmonella enterica* subspecies *enterica* serotype Enteritidis ($n = 170$) and *Salmonella enterica* subspecies *enterica* serotype Typhimurium ($n = 146$). These were followed by *Salmonella enterica* subspecies *enterica* serotype Anatum ($n = 62$) and *Salmonella enterica* subspecies *enterica* serotype Infantis ($n = 57$). *Salmonella enterica* subspecies *enterica* serotype Schwarzengrund and *Salmonella enterica* subspecies *enterica* serotype Muenchen were recovered in 50 and 48 cases, respectively. Of the total number of isolations recorded during the period under review, 871 (70.8%) occurred in poultry and other birds, 162 (13.2%) in horses, 116 (9.4%) in cattle, 26 (2.1%) in sheep and goats, 22 (1.8%) in rhinoceroses, 16 (1.3%) in pigs, 8 (0.6%) in crocodiles, 6 (0.5%) in cats and 6 (0.5%) in leopards. Food animals accounted for 83.5% of the total isolations, with cattle and poultry representing approximately 72.7%. Forty-two (3.4 %) isolates were found from non-food animals that include rhinoceroses ($n = 22$), crocodiles ($n = 8$), leopards ($n = 6$) and cats ($n = 6$). *Salmonella* Heidelberg was the most frequently isolated serotype, whereas *S. Typhimurium* had the widest zoological distribution. Clinical laboratory isolation of different *Salmonella* serotypes from various hosts may aid in recognising the threat to livestock, public and environmental health. Moreover, it may also highlight the potential zoonotic and food safety risk implications of the detected *Salmonella* serotypes.

Introduction

Salmonella infection (salmonellosis) is a common bacterial disease that mainly affects the intestinal tract of vertebrate and invertebrate hosts. *Salmonella* bacteria typically live in animal and human intestines and are shed in faeces. Salmonellosis remains a persistent threat to human and animal health and welfare in spite of the advances made in its detection, typing and control (Peek et al. 2004; Refsum et al. 2002). *Salmonella* and salmonellosis cause losses in the livestock and poultry industry because of death, abortion, decreased milk, meat and egg production, cost of testing and control programmes (Uzzau et al. 2000; Veling et al. 2002).

The genus *Salmonella* contains the species *Salmonella enterica*, *Salmonella bongori* and *Salmonella subterranean*, which was proposed in 2005 (Coburn, Grassl & Finlay 2007). However, according to the World Health Organization, the new species, *Salmonella subterranean*, does not belong to the genus *Salmonella* (Grimont & Weill 2007). *Salmonella enterica* subspecies are further classified into more than 2500 serovars or serotypes and include pathogens of great medical and veterinary importance (EFSA 2008; Mead et al. 1999; Morgan et al. 2004). These serovars differ greatly in their host range and degree of host adaptation. *Salmonella* serotypes are named in accordance with the Kaufmann–White typing system, defined by different combinations of somatic O, surface Vi and flagellar H antigens, which determine the serologically defined names appended to the *Salmonella* species as serovars or serotypes (Popoff 2001). In 2005, *Salmonella enterica* finally gained official approval as the type species of the genus *Salmonella* (Coburn et al. 2007).

The incidence of *Salmonella* infections in animals is closely associated with the age of the animals, husbandry and management – related factors in intensive farming that are conducive to the spread of infection resulting in an increase of clinical disease outbreaks (Baumler et al. 1998; Kidanemariam, Engelbrecht & Picard 2010). Clinically sick animals pose the greatest risk to

Read online:

Scan this QR code with your smart phone or mobile device to read online.

humans because they are more likely to shed *Salmonella* in higher concentrations than apparently healthy animals.

Salmonellosis is one of the most common food-borne diseases. High-protein foods such as meat, poultry, fish and eggs are most commonly associated with *Salmonella*. However, any food that becomes contaminated and held at temperatures that promote bacterial growth can cause salmonellosis (Altekruse et al. 2006; Kidanemariam et al. 2010; Kimura et al. 2004).

Periodic surveillance reports, such as this one, could provide useful information on the changing patterns of salmonellosis in animals and foods of animal origin. It also assists in the study of the epidemiology, risk factors associated with the disease, and control of the disease. The purpose of this report is to present the retrospective records of the animal isolations and the zoological distribution of the *Salmonella* serotypes isolated and serotyped at the Agricultural Research Council – Onderstepoort Veterinary Research (ARC-OVR) from 2007 to 2014.

Materials and methods

Specimens submitted to the ARC-OVR bacteriology laboratory comprised bacterial cultures, organs, tissues, swabs, faeces, fluid and tissues obtained after abortion and egg samples. Unprocessed and raw specimens were inoculated into buffered peptone water (pH 7.2) as pre-enrichment and incubated at 37 °C for 18–24 h. One millilitre of the pre-enrichment broth was transferred into 9 ml of Rappaport Vassiliadis (Oxoid®) enrichment broth and incubated at 42 °C for 18–24 h. Subcultures from enrichment media were grown on to selective solid media such as xylose-lysine deoxycholate (XLD) agar (Difco®) and incubated at 37 °C for 18–24 h. Black colonies with a pink periphery were considered presumptively positive for *Salmonella* and were further confirmed with biochemical tests. Only those Gram-negative isolates that were indole-negative, motile, Simmon's citrate-negative, urease negative, produced hydrogen sulphide in a triple sugar iron (TSI) slant, were lysine decarboxylase positive, fermented dulcitol but did not ferment lactose and were malonate negative were considered to be *Salmonella enterica*. Additional carbohydrate fermentation tests, such as gas production in Durham tubes and fermentation of sorbitol, arabinose, rhamnose, maltose and trehalose, were included to make provision for those *Salmonella* organisms that do not necessarily fit the above-mentioned criteria. Consideration was also given to *S. enterica* serovar Gallinarum and *S. enterica* serovar Pullorum that are non-motile when isolation and identification were done. There are variable reactions to Simmon's citrate utilisation and hydrogen production, and special attention was given to these features not to miss salmonellae.

Confirmed *Salmonella* isolates were further serotyped according to the Kauffmann–White classification scheme using a battery of somatic O and flagellar H polyvalent and monovalent antisera.

Data were captured in a dedicated Microsoft Excel™ data sheet for subsequent analysis. Descriptive statistics were employed to obtain values of proportions and percentiles.

Results

A total of 1229 *Salmonella* isolates were serotyped successfully during the surveillance period 2007–2014. The number of isolates and distribution of the various serotypes in different host species are shown in Table 1. The isolates were classified into 108 serotypes under the 17 groups of the Kauffmann–White classification scheme.

Of the total of 1229 isolates with 108 serotypes recorded during the period under review, 871 (70.8%) occurred in poultry and other birds, 162 (13.2%) in horses, 116 (9.4%) in cattle, 26 (2.1%) in sheep and goats, 22 (1.8%) in rhinoceroses, 16 (1.3%) in pigs, 8 (0.6%) in crocodiles, 6 (0.5%) in cats and 6 (0.5%) in leopards. Despite the large number of serotypes involved, the majority of isolates were limited to only a few serotypes. For example, from the total of 108 serotypes detected, only 8 serotypes contributed 63.0% (775 isolates), while 100 serotypes collectively accounted for 37.0% (454 isolates). The eight most common serotypes were *S. Anatum* (5.0%), *S. Dublin* (3.4%), *S. Enteritidis* (13.8%), *S. Heidelberg* (16.2%), *S. Infantis* (4.6%), *S. Muenchen* (3.9%), *S. Schwarzengrund* (4.0%) and *S. Typhimurium* (11.8%) (Table 1). The most frequently isolated serotype was *S. Heidelberg*, accounting for 16.2% of all isolates. Twenty-seven serotypes were isolated only once during the survey period. *S. Typhimurium* had the widest zoological distribution, followed by *S. Enteritidis* and *S. Heidelberg*.

In domestic fowl and other birds, the most common serotypes detected were *S. Enteritidis* (18.2%), *S. Heidelberg* (15.2%), *S. Typhimurium* (7.5%) and *S. Schwarzengrund* (5.4%); these four serotypes in total contributed 46.5% of the isolates, while the remaining 84 serotypes accounted for 53.5% of the total isolates (Tables 1 and 2).

In cattle, the most common serotypes detected were *S. Dublin* (33.6%), *S. Typhimurium* (14.6%), *S. Anatum* (11.2%) and *S. Infantis* (6.9%) and accounted for 66.4% of the total isolates, while the remaining 25 serotypes accounted for 33.6% of the total isolates (Table 3).

In horses, 162 isolations involving 18 serotypes were recorded in the period under review. The three major serotypes were *S. Heidelberg* (32.1%), *S. Anatum* (22.2%) and *S. Typhimurium* (21.6%). The remaining 15 serotypes collectively accounted for 24.8% of the total isolations (Tables 1 and 4).

Eight serotypes were identified from 26 isolations that occurred in sheep and goats. *Salmonella* Typhimurium (34.6%), *S. Enteritidis* (15.4%), *S. Infantis* (11.5%) and *S. Isangi* (11.5%) were the major serotypes and collectively accounted for 73.1% of the isolations (Table 5).

TABLE 1: Number of isolates and zoological distribution of the various serotypes 2007–2014.

Serotype	Cattle	Poultry	Leopard	Rhino	Crocodile	Feline	Equine	Sheep and goat	Pig	Total
S Aarhus	1	1	0	0	0	0	0	0	0	2
S Abaetetuba	0	7	0	0	0	0	0	0	0	7
S Aberdeen	3	9	0	0	0	0	0	0	0	12
S Achersleben	0	0	0	0	0	0	2	0	0	2
S Adeoyo	0	0	0	0	0	0	9	0	0	9
S Agona	0	12	0	0	0	0	0	2	0	14
S Alachua	0	3	0	0	0	0	0	0	0	3
S Anatum	13	13	0	0	0	0	36	0	0	62
S Berta	0	7	0	0	0	0	0	0	0	7
S Bovismorbificans	1	0	0	0	0	0	0	0	0	1
S Blockley	0	2	0	0	0	0	0	0	0	2
S Brandenburg	1	3	0	0	0	0	0	0	0	4
S Braenderup	0	6	0	0	0	0	0	0	0	6
S Brancaster	0	16	0	0	0	0	0	0	0	16
S Bredeney	1	3	0	0	0	0	0	0	0	4
S Brezany	0	1	0	0	0	0	0	0	0	1
S Canada	0	2	0	0	0	0	0	0	0	2
S Cardoner	6	0	0	0	0	0	0	0	0	6
S Cerro	0	1	0	0	0	0	0	0	0	1
S Chailey	0	2	0	0	0	0	0	0	0	2
S Chester	2	0	0	0	0	0	0	0	0	2
S Chicago	2	0	0	0	0	0	0	0	0	2
S Cremieu	1	0	0	0	0	0	0	0	0	1
S Claibornei	0	2	0	0	0	0	0	0	0	2
S Colorado	0	0	0	0	0	0	4	0	0	4
S Corvallis	0	4	0	0	0	0	0	0	0	4
S Cubana	1	0	0	0	0	0	0	0	0	1
S Derby	0	3	0	0	0	0	0	0	2	5
S Dublin	39	0	0	1	0	0	0	2	0	42
S Duesseldorf	1	2	0	0	0	0	0	0	0	3
S Duisburg	0	2	0	0	0	0	0	0	0	2
S Eastbourne	0	1	0	0	0	0	0	0	0	1
S Edinburg	0	2	0	0	0	0	0	0	0	2
S Eko	0	2	0	0	0	0	0	0	0	2
S Enteritidis	1	159	0	3	0	0	0	4	3	170
S Farmsen	0	6	0	0	0	0	0	0	0	6
S Fulda	0	4	0	0	8	0	0	0	0	12
S Give	0	0	0	0	0	0	0	0	1	1
S Glasgow	0	0	0	0	0	0	2	0	0	2
S Glostrup	0	1	0	0	0	0	0	0	0	1
S Gombe	0	1	0	0	0	0	0	0	0	1
S Goldcoast	2	0	0	0	0	0	0	0	0	2
S Gustavia	0	0	0	0	0	0	1	0	0	1
S Hadar	0	12	0	0	0	0	0	0	0	12
S Havana	0	5	0	0	0	0	0	0	0	5
S Hayindogo	0	4	0	0	0	0	0	0	0	4
S Heidelberg	1	133	0	12	0	2	52	0	0	200
S Idikan	0	3	0	0	0	0	0	0	0	3
S Indiana	0	7	0	0	0	0	0	0	0	7
S Infantis	8	41	0	0	0	0	5	3	0	57
S Isangi	0	3	0	0	0	0	0	3	0	6
S Israel	0	1	0	0	0	0	0	0	0	1
S Istanbul	0	1	0	0	0	0	0	0	0	1
S Jerusalem	0	2	0	0	0	0	0	0	0	2
S Kainji	0	2	0	0	0	0	0	0	0	2
S Kalamu	0	1	0	0	0	0	2	0	0	3
S Kedougou	2	0	0	0	0	0	0	0	0	2
S Kentucky	0	7	0	0	0	0	0	0	0	7
S Kiambu	1	4	0	0	0	0	0	0	0	5
S Kingston	0	1	0	0	0	0	0	0	0	1
S Kotu	0	3	0	0	0	0	0	0	0	3

Table 1 continues on the next page →

TABLE 1 (Continues...): Number of isolates and zoological distribution of the various serotypes 2007–2014.

Serotype	Cattle	Poultry	Leopard	Rhino	Crocodile	Feline	Equine	Sheep and goat	Pig	Total
S Kottbus	1	0	0	0	0	0	0	0	0	1
S Kouka	0	2	0	0	0	0	0	0	0	2
S Lexington	0	1	0	0	0	0	0	0	0	1
S Lockleaze	0	2	0	0	0	0	0	0	0	2
S London	0	4	0	0	0	0	0	0	0	4
S Mandera	0	0	0	0	0	0	1	0	0	1
S Manhattan	0	1	0	0	0	0	2	0	0	3
S Mbandaka	0	6	0	0	0	0	0	0	0	6
S Minnesota	0	25	0	0	0	0	0	0	0	25
S Mikawasima	0	0	6	0	0	0	0	0	0	6
S Montevideo	0	10	0	0	0	0	0	0	0	10
S Muenchen	2	43	0	0	0	0	2	1	0	48
S Newport	0	22	0	0	0	0	0	0	0	22
S Ohio	0	3	0	0	0	0	0	0	0	3
S Orion	1	6	0	0	0	0	0	0	0	7
S Oslo	0	1	0	0	0	0	0	0	0	1
S Othmarschen	1	2	0	0	0	0	0	0	0	3
S Paratyphi	0	5	0	0	0	0	0	0	0	5
S Planckendael	0	1	0	0	0	0	0	0	0	1
S Poitiers	0	1	0	0	0	0	0	0	0	1
S Panama	0	1	0	0	0	0	0	0	0	1
S Poona	0	1	0	0	0	0	0	0	0	1
S Pretoria	2	3	0	0	0	0	2	0	0	7
S Reading	2	2	0	0	0	0	0	0	0	4
S Rideau	0	0	0	0	0	0	1	0	0	1
S Rissen	0	5	0	0	0	0	0	0	0	5
S Ruiru	0	4	0	0	0	0	0	0	0	4
S Saintpaul	0	11	0	0	0	0	0	0	0	11
S Salford	1	0	0	0	0	0	0	0	0	1
Sambre	0	0	0	0	0	0	1	0	0	1
S Sandiego	0	0	0	0	0	0	0	0	1	1
S Schoeneberg	1	0	0	0	0	0	2	0	0	3
S Schwarzengrund	0	47	0	0	0	0	3	0	0	50
S Senftenberg	0	25	0	0	0	0	0	0	0	25
S Shubra	0	2	0	0	0	0	0	0	0	2
S Stanley	0	1	0	0	0	0	0	0	0	1
S Stanleyville	0	13	0	0	0	0	0	0	0	13
S Stratford	1	0	0	0	0	0	0	0	0	1
S Tallahassee	0	2	0	0	0	0	0	0	0	2
S Tennessee	0	7	0	0	0	0	0	0	0	7
S Thompson	0	13	0	0	0	0	0	0	0	13
S Tsevie	0	2	0	0	0	0	0	0	0	2
S Typhimurium	17	66	0	6	0	4	35	9	9	146
S Virchow	0	13	0	0	0	0	0	0	0	13
S Winneba	0	6	0	0	0	0	0	0	0	6
S Wippra	0	4	0	0	0	0	0	0	0	4
S Yoruba	0	1	0	0	0	0	0	2	0	3
Total	116	871	6	22	8	6	162	26	16	1233

Sixteen isolations involving five serotypes were recorded in pigs. The majority of isolates were *S. Typhimurium*, that accounted for 56.3% of the cases, followed by *S. Enteritidis* (18.7%) and *S. Derby* (12.5%) (Table 6). In captive and wild animals, 42 isolates involving 6 serotypes were recorded (Table 7). *Salmonella* Heidelberg (33.33%) and *S. Typhimurium* (23.8%) were the two major serotypes isolated during the survey period. *Salmonella* Fulda (19.0%) and *S. Mikawasima* (14.3%) were isolated exclusively from crocodiles and leopards, respectively (Table 7). *Salmonella* Dublin (2.4%) and *S. Enteritidis* (7.1%) were detected from rhinoceroses.

Discussion

Laboratory diagnosis continues to provide important epidemiological information that contributes significantly to continuous disease surveillance programmes in the country, if properly validated tests are used. The data set presented in this report is a follow-up of similar laboratory surveillance information carried out from 1996 through 2006 as described elsewhere (Kidanemariam et al. 2010).

There has been a steady increase over the years in the number of *Salmonella* serotypes isolated and in the number of animal

TABLE 2: Common *Salmonella* serotypes in poultry in South Africa.

Isolate	2007	2008	2009	2010	2011	2012	2013	2014	Total
S Enteritidis									
Count	26	14	10	62	2	12	31	2	159
%	16.88	14.00	10.30	42.42	6.89	7.69	19.25	8.33	18.25
S Heidelberg									
Count	20	0	11	5	0	63	34	0	133
%	12.98	0.00	11.34	3.42	0.00	40.38	21.11	0.00	15.26
S Infantis									
Count	1	6	5	4	8	3	8	6	41
%	0.64	6.00	5.15	2.73	27.58	1.92	4.96	25.00	4.70
S Minnesota									
Count	2	6	1	3	0	4	9	0	25
%	1.29	6.00	1.03	2.05	0.00	2.56	5.59	0.00	2.87
S Muenchen									
Count	2	15	0	2	0	17	7	0	43
%	1.29	15.00	0.00	1.36	0.00	10.89	4.34	0.00	4.93
S Newport									
Count	0	5	3	0	3	5	3	3	22
%	0.00	5.00	3.09	0.00	10.34	3.20	1.86	12.50	2.52
S Schwarzengrund									
Count	0	5	2	8	2	5	18	7	47
%	0.00	5.00	2.06	5.47	6.89	3.20	11.18	29.16	5.39
S Senftenberg									
Count	3	1	15		5	0	1	0	25
%	1.94	1.00	15.46	0.00	17.24	0.00	0.62	0.00	2.87
S Typhimurium									
Count	22	8	20	18	1	9	7	0	66
%	14.28	8.00	20.61	12.32	3.44	5.76	4.34	0.00	7.57
Others									
Count	78	48	34	44	16	41	43	6	310
%	50.64	48.00	35.05	30.13	55.17	26.28	26.70	25.00	35.59
Total									
Count	154	100	97	146	29	156	161	24	871
%	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00

TABLE 3: Common *Salmonella* serotypes in cattle in South Africa.

Isolate	2007	2008	2009	2010	2011	2012	2013	2014	Total
S Anatum									
Count	1	5	1	0	0	5	1	0	13
%	5.00	27.77	6.66	0.00	0.00	33.33	4.00	0.00	11.20
S Dublin									
Count	8	2	6	9	3	2	6	3	39
%	40.00	11.11	40.00	60.00	75.00	13.33	24.00	75.00	33.60
S Infantis									
Count	1	2	0	0	0	2	3	0	8
%	5.00	11.11	0.00	0.00	0.00	13.33	12.00	0.00	6.89
S Typhimurium									
Count	7	4	1	1	0	3	1	0	17
%	35.00	22.22	6.66	6.66	0.00	20.00	4.00	0.00	14.65
Others									
Count	3	5	7	5	1	3	14	1	39
%	15.00	27.77	46.66	33.30	25.00	20.00	56.00	25.00	33.62
Total									
Count	20	18	15	15	4	15	25	4	116
%	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00

species involved. *Salmonella* serotypes clearly seem to differ in their distributions between different animal populations (Table 1). For instance, approximately 95.3% of all detected *Salmonella* serotypes are predominantly associated with domestic species, namely poultry, cattle, horses and sheep and goats, whereas rhinoceroses, leopards, crocodiles and

pigs contributed only 4.7%. However, care must be taken when considering these data because of the possibility of a skewed focus from targeted sampling of mostly food animals as compared to wild and game animals. In spite of this, the data probably give a reasonable assessment of the incidence of salmonellosis and the *Salmonella* serotypes involved.

TABLE 4: Common *Salmonella* serotypes in horses in South Africa.†

Isolate	2008	2009	2011	2012	2013	2014	Total
S Achersleben							
Count	0	0	1	0	0	1	2
%	0.00	0.00	1.72	0.00	0.00	2.12	1.23
S Adeoyo							
Count	0	0	0	0	9	0	9
%	0.00	0.00	0.00	0.00	20.45	0.00	5.56
S Anatum							
Count	4	0	8	4	12	8	36
%	100.00	0.00	13.79	100.00	27.27	17.02	22.22
S Colorado							
Count	0	0	2	0	0	2	4
%	0.00	0.00	3.45	0.00	0.00	4.25	2.47
S Glasgow							
Count	0	0	0	0	2	0	2
%	0.00	0.00	0.00	0.00	4.54	0.00	1.23
S Gustaria							
Count	0	0	0	0	1	0	1
%	0.00	0.00	0.00	0.00	2.27	0.00	0.62
S Heidelberg							
Count	0	0	32	0	0	20	52
%	0.00	0.00	55.17	0.00	0.00	42.55	32.1
S Infantis							
Count	0	0	0	0	5	0	5
%	0.00	0.00	0.00	0.00	11.36	0.00	3.08
S Kalamu							
Count	0	0	1	0	0	2	2
%	0.00	0.00	1.72	0.00	0.00	4.25	1.23
S Mandera							
Count	0	0	0	0	1	0	1
%	0.00	0.00	0.00	0.00	2.27	0.00	0.61
S Mbandaka							
Count	0	0	1	0	0	1	2
%	0.00	0.00	1.72	0.00	0.00	2.12	1.23
S Muenchen							
Count	0	0	1	0	0	1	2
%	0.00	0.00	1.72	0.00	0.00	2.12	1.23
S Pretoria							
Count	0	0	0	0	2	0	2
%	0.00	0.00	0.00	0.00	4.54	0.00	1.23
S Rideau							
Count	0	0	0	0	1	0	1
%	0.00	0.00	0.00	0.00	2.27	0.00	0.61
S Sambre							
Count	0	0	0	0	1	0	1
%	0.00	0.00	0.00	0.00	2.27	0.00	0.61
S Schoeneberg							
Count	0	0	1	0	0	1	2
%	0.00	0.00	1.72	0.00	0.00	2.12	1.23
S Schwarzengrund							
Count	0	1	0	0	2	0	3
%	0.00	16.70	0.00	0.00	4.54	0.00	1.85
S Typhimurium							
Count	0	5	11	0	8	11	35
%	0.00	83.33	18.96	0.00	18.18	23.40	21.60
Total							
Count	4	6	58	4	44	47	162
%	100.00	100.00	100.00	100.00	100.00	100.00	100.00

†, No samples were presented for testing between 2007 and 2010.

Over 2500 *Salmonella* serovars are recognised worldwide (Coburn et al. 2007; EFSA 2008), and the number continues to rise. However, despite the existence of formidable number

of different serotypes, only a few are commonly associated with clinical disease in humans and animals (Warnick et al. 2001). It is also noted in the current study that despite the

TABLE 5: Common *Salmonella* serotypes in sheep and goats in South Africa.

Isolate	2010	2012	2013	Total
S Agona				
Count	0	1	1	2
%	0.00	14.28	7.69	7.69
S Dublin				
Count	0	0	2	2
%	0.00	0.00	15.38	7.69
S Enteritidis				
Count	1	1	2	4
%	16.66	14.28	15.38	15.38
S Infantis				
Count	0	0	3	3
%	0.00	0.00	23.07	11.53
S Isangi				
Count	1	2	0	3
%	16.66	28.57	0.00	11.53
S Muenchen				
Count	0	1	0	1
%	0.00	14.28	0.00	3.84
S Typhimurium				
Count	3	1	5	9
%	50.00	14.28	38.46	34.61
S Yoruba				
Count	1	1	0	2
%	16.66	14.28	0.00	7.69
Total				
Count	6	7	13	26
%	100.00	100.00	100.00	100.00

TABLE 6: Common *Salmonella* serotypes in pigs in South Africa.

Isolate	2008	2009	2010	2011	2012	2013	2014	Total
S Derby								
Count	1	0	0	0	1	0	0	2
%	20.00	0.00	0.00	0.00	20.00	0.00	0.00	12.50
S Enteritidis								
Count	1	0	0	0	1	1	0	3
%	20.00	0.00	0.00	0.00	20.00	25.00	0.00	18.75
S Give								
Count	0	0	0	0	0	1	0	1
%	0.00	0.00	0.00	0.00	0.00	25.00	0.00	6.25
S Sandiego								
Count	0	0	1	0	0	0	0	1
%	0.00	0.00	100.00	0.00	0.00	0.00	0.00	6.25
S Typhimurium								
Count	3	0	0	0	3	2	1	9
%	60.00	0.00	0.00	0.00	60.00	50.00	100.00	56.25
Total								
Count	5	0	1	0	5	4	1	16
%	100.00	0.00	100.00	0.00	100.00	100.00	100.00	100.00

large number of serotypes involved, the majority of isolates were mainly represented by a small number of serotypes. For instance, from the total of 108 serotypes detected among 1229 isolates, only 8 serotypes contributed 63.0 % (775 isolates), while 100 serotypes collectively accounted for 37.0% (454 isolates).

Salmonella serotypes can be divided into host-specific, host-adapted and generalist serotypes, with important implications for the epidemiology and risk factors of the diseases in the associated host species (Baumler et al. 1998; Uzzau et al. 2000).

Host-specific serotypes, such as *S. Paratyphi A* in humans and *S. Gallinarum* biovars *Gallinarum* and *Pullorum* in chickens, only caused disease in one host species (Pascopella et al. 1995; Uzzau et al. 2000). These host-specific serotypes were not isolated during the survey period. *Salmonella Paratyphi A* is exclusively a human pathogen, and samples are submitted to public health facilities. Nonetheless, *S. Typhi* was confirmed through blood and stool cultures from cases of a typhoid fever outbreak in South Africa (Anon 2016). The absence of *S. Gallinarum* and *S. Pullorum* in the data set of the current survey could be because of the regular targeted

TABLE 7: Common *Salmonella* serotypes in wild and captive animals in South Africa.†

Isolate	2011	2012	2013	2014	Total
S Dublin					
Count	0	0	0	1	1
%	0.00	0.00	0.00	4.76	2.38
S Enteritidis					
Count	1	0	1	1	3
%	5.00	0.00	100.00	4.76	7.14
S Fulda					
Count	4	0	0	4	8
%	20.00	0.00	0.00	19.04	19.04
S Mikawasima					
Count	3	0	0	3	6
%	15.00	0.00	0.00	14.28	14.28
S Heidelberg					
Count	7	0	0	7	14
%	35.00	0.00	0.00	33.33	33.33
S Typhimurium					
Count	5	0	0	5	10
%	25.00	0.00	0.00	23.81	23.81
Total					
Count	20	0	1	21	42
%	100.00	0.00	100.00	100.00	100.00

†, No samples were presented for testing between 2007 and 2010.




FIGURE 1: Annual *Salmonella* isolations and number of serotypes from animal origin.

monitoring programme that assisted poultry farms to control and prevent *Salmonella* infections in general and infections as a result of *S. Gallinarum* and *S. Pullorum* in particular (A.K. Gelaw, unpublished results). In addition, small incidences are less likely to be reported and to be passed unnoticed. Furthermore, the lack of either serovar in the current survey may also be attributed to the selection criteria in the methodology that often targets motile salmonellae.

Host-adapted serotypes are predominantly associated with a single host species, but reports suggest that these serotypes can cause disease in other species as well (Pascopella et al. 1995; Uzzau et al. 2000). For example, *S. Dublin*, *S. Enteritidis* and *S. Choleraesuis* were previously regarded as host-adapted serotypes for cattle, chicken and pigs, respectively (Anderson et al. 1997; Uzzau et al. 2000). However, it became apparent that these serotypes commonly cause disease in a broad range of hosts and may be considered as generalist serotypes. This is supported by the results of the current survey where *S. Dublin* was isolated from cattle, goats and

rhinoceroses, and *S. Enteritidis* was isolated from poultry, pigs, sheep and rhinoceroses, demonstrating the ability of these serotypes to adapt and infect multiple hosts.

Generalist serotypes such as *Salmonella* Typhimurium, on the other hand, are associated with a wide range of hosts. The current data supported this assertion, as *S. Typhimurium* was detected across a wide range of animal species that include poultry, cattle, horses, sheep, goats and pigs (Table 1). This serotype was also found in wild and captive animals such as a leopard, rhinoceros and crocodile. Similar studies have demonstrated that *S. Typhimurium* is the most common serovar isolated in livestock (Morgan et al. 2004; Rabsch et al. 2002) and human non-typhoidal infections, especially in immune-compromised patients (Calvert, Stewart & Reilly 1998; Gordon 2008; Keddy et al. 2017; Voetsch et al. 2004).

Salmonella Heidelberg is one of the generalist serotypes and was by far the most common serotype, representing 16.2% of the total isolations, with the majority originating from horses, in which species it accounted for 32.1% of the total cases. This result could partially be attributed to the increased epidemiological monitoring to ascertain the causes of clinical episodes of enteric infections in the hospitalised horses at the Veterinary Academic Hospital of the University of Pretoria (J. Gouws, pers. com., June 2015). Faecal material was submitted for testing, and *S. Heidelberg* was isolated from horses admitted to the hospital. Because of the potential risk of spread of *Salmonella* from an infected horse, the equine hospital has introduced biosecurity measures to help prevent disease transmission to personnel, the environment and other patients. Similar reports showed that hospitalised horses were more likely to shed *Salmonella* than horses housed in their own barns, possibly because of stress or

illness (Amavisit et al. 2001). A study at a large veterinary hospital in the United States demonstrated that 13% of horses admitted for colic were shedding *Salmonella* species (Kim et al. 2001).

Salmonella Typhimurium exhibited the widest zoological distribution (Table 1). This finding is not unexpected because *S. Typhimurium* has been thought of as the most ubiquitous and broadest host-range serotype, as it is frequently associated with diseases in numerous host species, including humans, livestock, domestic fowl, rodents and birds (Bahnsen et al. 2006; Padungtod, Kadohira & Hill 2008). It was, however, demonstrated in the current survey that the incidence of *S. Typhimurium* was relatively higher in poultry and equines with incidence rates of 45.2% and 23.9%, respectively.

Although *S. Enteritidis* was isolated from various host species, it was by far the most common serotype encountered in poultry, accounting for 93.5% of the incidents. The remaining 6.5% were shared among cattle (0.6%), rhinoceros (1.7%), sheep (2.3%) and pigs (1.7%). This serotype is among the most common pathogens of chicken that could also adversely affect the health of human beings and other animal species following exposures (Kidaneariam et al. 2010). Vertical transmission with internal transovarian contamination of egg-yolk with *S. Enteritidis* has been confirmed, making uncooked eggs no longer safe to eat (Altekruse et al. 1993). Furthermore, *S. Enteritidis* was confirmed as causes of human outbreaks of non-typhoidal salmonellosis associated with the consumption of foods of animal origin in South Africa (Muvhali et al. 2017). *Salmonella* Enteritidis is often presented separately from other serotypes of *Salmonella* because this bacterium is specifically cited in zoonosis control legislation. The South African government has included *S. Enteritidis* in the list of controlled diseases (Anon 1984; Kidaneariam et al. 2010).

Salmonella Dublin and *Salmonella* Typhimurium are the two predominant serotypes detected in cattle in the current survey, accounting for 48.3% of the total recorded isolations. The relative incidence of *S. Dublin* (33.6%) is, however, higher than that of *S. Typhimurium* (14.6%) for the survey period. Similar studies have shown that *S. Dublin* may potentially be the most frequently isolated serotype in cattle, more than the broad host-range serotypes, such as *S. Typhimurium* (Brackelsberg, Nolan & Brown 1997; Kidaneariam et al. 2010). It should be noted that the disease epidemiology of *S. Dublin* and *S. Typhimurium* varies considerably. Most importantly, cattle infected with *S. Dublin* often become asymptomatic carriers and continue to excrete large numbers of organisms in their faeces for many years and often for life (Giles, Hopper & Wray 1989; Rice, Besser & Hancock 1997; Vanselow et al. 2007). Effective control of infection should thus include removal of chronically infected cattle and implementation of stringent biosecurity measures. On the other hand, hosts infected with *S. Typhimurium* will only shed the organism for a few weeks

to a month after clinical recovery (Huston, Wittum & Love 2002). However, it should be noted that *S. Typhimurium* tends to persist in the environment for longer periods (Rabsch et al. 2002). One study estimated the median duration of shedding in dairy cattle to equal 50 days, with a maximum duration of 391 days (Cummings et al. 2009). This can be mitigated by implementing proper biosecurity measures at all times.

Clinically affected herds and certain management systems may pose an increased risk to public health. Little is known about the *Salmonella* risk posed to humans by indirect animal contact, especially through environmental contamination. Considering the public health significance of this group of bacteria, further studies, especially on spatial clustering of human cases around livestock premises, are needed to assess the indirect risks posed by livestock operations in South Africa.

Acknowledgements

This work was supported partly by a grant from the Agricultural Research Council – Onderstepoort Veterinary Research (P10000046). We acknowledge Mrs Maluleka for the technical support during data retrieval. We are also indebted to Dr Sabeta for his critical comments on the manuscript.

Competing interests

The authors declare that there is no conflict of interest with regard to the writing of this article.

Authors' contributions

A.K.G. contributed to the study design, interpretation of data and preparation of the manuscript. I.M. and P.N. contributed to laboratory testing and data collection.

References

- Altekruse, S.F., Bauer, N., Chanlongbutra, A., DeSagun, R., Naugle, A., Schlosser, W. et al., 2006, '*Salmonella* Enteritidis in broiler chickens, United States, 2000–2005', *Emerging Infectious Diseases* 12(12), 1848–1852. <https://doi.org/10.3201/eid1212.060653>
- Altekruse, S., Koehler, J., Hickman-Brenner, F., Tauxe, R.V. & Ferris, K., 1993, 'A comparison of *Salmonella* Enteritidis phage types from egg-associated outbreaks and implicated laying flocks', *Epidemiology and Infection* 110, 17–22. <https://doi.org/10.1017/S0950268800050639>
- Amavisit, P., Markham, P.F., Lightfoot, D., Whithear, K.G. & Browning, G.F., 2001, 'Molecular epidemiology of *Salmonella* Heidelberg in an equine hospital', *Veterinary Microbiology* 80, 85–98.
- Anderson, R.L., Walker, R.L., Hird, D.W. & Blanchard, P.C., 1997, 'Case-control study of an outbreak of clinical disease attributable to *Salmonella* infection in eight dairy herds', *Journal of the American Veterinary Medical Association* 210, 528–530. [https://doi.org/10.1016/S0378-1135\(00\)00373-4](https://doi.org/10.1016/S0378-1135(00)00373-4)
- Bahnsen, P.B., Fedorka-Cray, P.J., Ladely, S.R. & Mateus-Pinilla, N.E., 2006, 'Herd-level risk factors for *Salmonella enterica* subsp. *enterica* in U.S. market pigs', *Preventive Veterinary Medicine* 76(3–4), 249–262. <https://doi.org/10.1016/j.prevetmed.2006.05.009>
- Baumler, A.J., Tsois, R.M., Ficht, T.A. & Adams, L.G., 1998, 'Evolution of host adaptation in *Salmonella enterica*', *Infection and Immunity* 66(10), 4579–4587.
- Brackelsberg, C.A., Nolan, L.K. & Brown, J., 1997, 'Characterization of *Salmonella* Dublin and *Salmonella* Typhimurium (Copenhagen) isolates from cattle', *Veterinary Research Communications* 21(6), 409–420. <https://doi.org/10.1023/A:1005803301827>
- Calvert, N., Stewart, W.C. & Reilly, W.J., 1998, '*Salmonella* Typhimurium DT104 infection in people and animals in Scotland: A collaborative epidemiological study 1993–96', *Veterinary Record* 143(13), 351–354. <https://doi.org/10.1136/vr.143.13.351>

- Coburn, B., Grassl, G.A. & Finlay, B.B., 2007, 'Salmonella, the host and disease: A brief review', *Immunology and Cell Biology* 85, 112–118. <https://doi.org/10.1038/sj.icb.7100007>
- Cummings, K.J., Warnick, L.D., Alexander, K.A., Cripps, C.J., Grohn, Y.T., James, K.L. et al., 2009, 'The duration of fecal *Salmonella* shedding following clinical disease among dairy cattle in the northeastern USA', *Preventive Veterinary Medicine* 92, 134–139. <https://doi.org/10.1016/j.prevetmed.2009.07.002>
- Department of Agriculture, Forestry & Fisheries, 1984, *Animal Diseases Act 1984 (Act 35 of 1984)*, Republic of South Africa.
- EFSA (European Food Safety Authority), 2008, 'Scientific opinion of the panel on biological hazards on a request from the European Commission on a quantitative microbiological risk assessment on *Salmonella* in meat: Source attribution for human salmonellosis from meat', *EFSA Journal* 625, 1–32.
- Giles, N., Hopper, S.A. & Wray, C., 1989, 'Persistence of *S. Typhimurium* in a large dairy herd', *Epidemiology and Infection* 103(2), 235–241. <https://doi.org/10.1017/S0950268800030582>
- Gordon, M., 2008, '*Salmonella* infections in immunocompromised adults', *Journal of Infections* 56, 413–422. <https://doi.org/10.1016/j.jinf.2008.03.012>
- Grimont, P.A. & Weill, F.-X., 2007, *Antigenic formulae of the Salmonella Serovars*, 9th edn., WHO Collaborating Centre for Reference and Research on Salmonella, Institut Pasteur, Paris.
- Huston, C.L., Wittum, T.E. & Love, B.C., 2002, 'Persistent fecal *Salmonella* shedding in five dairy herds', *Journal of the American Veterinary Medical Association* 220(5), 650–655. <https://doi.org/10.2460/javma.2002.220.650>
- Keddy, K.H., Takuva, S., Musekiwa, A., Puren, A.J., Sooka, A., Karstaedt, A. et al., 2017, 'An association between decreasing incidence of invasive non-typhoidal salmonellosis and increased use of antiretroviral therapy, Gauteng Province, South Africa, 2003–2013', *PLoS One* 12(3), e0173091. <https://doi.org/10.1371/journal.pone.0173091>
- Kidanemariam, A., Engelbrecht, M. & Picard, J., 2010, 'Retrospective study on the incidence of *Salmonella* isolations in animals in South Africa, 1996 to 2006', *Journal of the South African Veterinary Association* 81(1), 37–44. <https://doi.org/10.4102/jsava.v81i1.94>
- Kim, L.M., Morley, P.S., Traub-Dargatz, J.L., Salman, M.D. & Gentry-Weeks, C., 2001, 'Factors associated with *Salmonella* shedding among equine colic patients at a veterinary teaching hospital', *Journal of the American Veterinary Medical Association* 218, 740–748. <https://doi.org/10.2460/javma.2001.218.740>
- Kimura, A.C., Reddy, V., Marcus, R., Cieslak, P.R., Mohle-Boetani, J.C. & Kassenborg, H.D., 2004, 'Chicken consumption is a newly identified risk factor for sporadic *Salmonella enterica* serotype Enteritidis infections in the United States: A case-control study in FoodNet sites', *Clinical Infectious Diseases* 38, 244–252. <https://doi.org/10.1086/381576>
- Mead, P.S., Slutsker, L., Dietz, V., McCaig, L.F., Bresee, J.S., Shapiro, C. et al., 1999, 'Food-related illness and death in the United States', *Emerging Infectious Diseases* 5(5), 607–625. <https://doi.org/10.3201/eid0505.990502>
- Morgan, E., Campbell, J.D., Rowe, S.C., Bispham, J., Stevens, M.P., Bowen, A.J. et al., 2004, 'Identification of host-specific colonization factors of *Salmonella enterica* serovar Typhimurium', *Molecular Microbiology* 54, 994–1010. <https://doi.org/10.1111/j.1365-2958.2004.04323.x>
- Muvhali, M., Smith, A.M., Rakgantsu, A.M. & Keddy, K.H., 2017, 'Investigation of *Salmonella* Enteritidis outbreaks in South Africa using multi-locus variable number tandem repeats analysis, 2013–2015', *BMC Infectious Diseases* 17(1), 661. <https://doi.org/10.1186/s12879-017-2751-8>
- National Institute of Communicable Diseases, Communicable Diseases Communiqué, 2016, '*Salmonella* Typhi cases in South Africa, 2016', vol. 15(2), p. 1, Sandringham, Republic of South Africa.
- Padungtod, P., Kadohira, M. & Hill, G., 2008, 'Livestock production and foodborne diseases from food animals in Thailand', *Journal of Veterinary Medical Science* 70(9), 873–879. <https://doi.org/10.1292/jvms.70.873>
- Pascopella, L., Raupach, B., Ghori, N., Monack, D., Falkow, S. & Small, P.L., 1995, 'Host restriction phenotypes of *Salmonella* Typhi and *Salmonella* Gallinarum', *Infection and Immunity* 63, 4329–4335.
- Peek, S.F., Hartmann, F.A., Thomas, C.B. & Nordlund, K.V., 2004, 'Isolation of *Salmonella* spp from the environment of dairies without any history of clinical salmonellosis', *Journal of the American Veterinary Medical Association* 225(4), 574–577. <https://doi.org/10.2460/javma.2004.225.574>
- Popoff, M.Y., 2001, *Antigenic formulae of the Salmonella serovars*, 8th edn., WHO Collaborating Centre for Reference and Research on Salmonella, Institut Pasteur, Paris.
- Rabsch, W., Andrewes, H.L., Kingsley, R.A., Prager, R., Tschäpe, H., Adams, L.G. et al., 2002, '*Salmonella enterica* serotype Typhimurium and its host adapted variants', *Infection and Immunity* 70, 2249–2255. <https://doi.org/10.1128/IAI.70.5.2249-2255.2002>
- Refsum, T., Heir, E., Kapperud, G., Vardund, T. & Holstad, G., 2002, 'Molecular epidemiology of *Salmonella enterica* serovar Typhimurium isolates determined by pulsed-field gel electrophoresis: Comparison of isolates from avian wildlife, domestic animals, and the environment in Norway', *Applied and Environmental Microbiology* 68(11), 5600–5606. <https://doi.org/10.1128/AEM.68.11.5600-5606.2002>
- Rice, D.H., Besser, T.E. & Hancock, D.D., 1997, 'Epidemiology and virulence assessment of *Salmonella* Dublin', *Veterinary Microbiology* 56, 111–124. [https://doi.org/10.1016/S0378-1135\(96\)01352-1](https://doi.org/10.1016/S0378-1135(96)01352-1)
- Uzzau, S., Brown, D.J., Wallis, T., Rubino, S., Leori, G., Bernard, S. et al., 2000, 'Host adapted serotypes of *Salmonella enterica*', *Epidemiology and Infection* 125(2), 229–255. <https://doi.org/10.1017/S0950268899004379>
- Vanselow, B.A., Hum, S., Hornitzky, M.A., Eamens, G.J. & Quinn, K., 2007, '*Salmonella* Typhimurium persistence in a Hunter Valley dairy herd', *Australian Veterinary Journal* 85(11), 446–450. <https://doi.org/10.1111/j.1751-0813.2007.00224.x>
- Veling, J., Wilpshaar, H., Frankena, K., Bartels, C. & Barkema, H., 2002, 'Risk factors for clinical *Salmonella enterica* subsp. *enterica* serovar Typhimurium infection on Dutch dairy farms', *Preventive Veterinary Medicine* 54(2), 157–168. [https://doi.org/10.1016/S0167-5877\(02\)00023-5](https://doi.org/10.1016/S0167-5877(02)00023-5)
- Voetsch, A.C., Van Gilder, T.J., Angulo, F.J., Farley, M.M., Shallow, S., Marcus, R. et al., 2004, 'FoodNet estimate of the burden of illness caused by non-typhoidal *Salmonella* infections in the United States', *Clinical Infectious Disease* 38, 127–134. <https://doi.org/10.1086/381578>
- Warnick, L.D., Crofton, L.M., Pelzer, K.D. & Hawkins, M.J., 2001, 'Risk factors for clinical salmonellosis in Virginia, USA, cattle herds', *Preventive Veterinary Medicine* 49, 259–275. [https://doi.org/10.1016/S0167-5877\(01\)00172-6](https://doi.org/10.1016/S0167-5877(01)00172-6)