

The effect of dietary protein on reproduction in the mare. V. Endocrine changes and conception during the early *post partum* period

F E van Niekerk^a and C H van Niekerk^a

ABSTRACT

Pregnant Anglo-Arab and Thoroughbred mares ($n = 24$) were divided randomly according to age and breed into 4 groups of 6 mares each from approximately 6 weeks before their expected foaling date. Diets received by the 4 groups varied in essential amino-acid and total protein contents. Serum progesterone, FSH and LH concentrations were determined from the day of parturition until foal heat and during the 1st oestrous cycle following foal heat. Serum progesterone, FSH and LH concentrations did not differ between the treatment groups. Progesterone concentrations were high ($\bar{x} = 7.0$: 5.2–16.4 ng/ml) at parturition but decreased rapidly within 48 h. As progesterone concentrations decreased LH concentrations increased from Days 3–6 *post partum* to reach maximum values at, or the day after ovulation. FSH concentrations declined 3–4 d after parturition and increased 2–3 d before ovulation at foal heat. The duration of elevated progesterone concentrations during the luteal phase of the subsequent oestrous cycle affected the interovulatory period. A 12–14 d FSH cyclical releasing pattern occurred. Season/photoperiod affected the resumption of normal oestrous cyclicity during the *post partum* period. The duration of the 1st oestrous cycle after foal heat in mares fed a low-quality protein diet showed a greater range (13–30 d) compared to mares fed a high-quality protein diet (18–26 d).

Key words: equine, FSH, LH, *post partum* period, protein nutrition, serum progesterones.

Van Niekerk F E, Van Niekerk C H The effect of dietary protein on reproduction in the mare. V. Endocrine changes and conception during the early *post partum* period. *Journal of the South African Veterinary Association* (1998) 69(3): 81–88 (En.). Department of Human and Animal Physiology, University of Stellenbosch, Private Bag X1, Matieland, 7602 South Africa.

INTRODUCTION

The duration of pregnancy of *c.* 11 months in the Thoroughbred mare implies that the mare must reconceive within 30 days *post partum* in order to produce a foal each year. The *post partum* interval is described as the time after parturition when the uterus undergoes involution to the state where pregnancy can be maintained¹⁴. Late pregnancy, parturition and the early *post partum* period are characterised by major changes in the reproductive and endocrine systems³².

Foal heat occurs within 7–9 d of parturition in more than 90 % of mares^{4,8,19,31}. Although mares will ovulate and conceive during foal heat¹⁶, several studies indicated a reduced pregnancy rate^{4,28} and embryonic loss that may be associated with the actual state of the uterine environment¹⁴.

The reproductive efficiency of lactating mares is affected by both season/photo-period and nutrition. Approximately 28 % of Thoroughbred mares that foal early in spring do not show signs of oestrus³⁰. Optimal energy intake of lactating mares is important because this is the period when reconception should take place⁷. The bodily condition of the mare at foaling is therefore important, as mares in poor condition at foaling may maintain that condition at the expense of milk production¹⁷. Conception is improved when mares are maintained in a positive energy balance during lactation^{25,37} and by limiting the period of suckling by the foal¹⁶. Studies have indicated that a protein deficiency in the diet might contribute to early embryonic loss irrespective of the mare's bodily condition^{5,38}.

The purpose of this study was to determine the normal clinical, morphological and endocrine changes that occur in the reproductive functions of the mare during the *post partum* period until reconception. The effects of season at the

time of foaling and dietary protein quality and intake were also investigated.

MATERIALS AND METHODS

Pregnant Thoroughbred and Anglo-Arab mares ($n = 24$) were kept on pasture until 6 weeks before their expected foaling dates and were then allocated randomly according to age and breed to 1 of 4 dietary groups of 6 mares each. The 4 diets consisted of: Group 1: tef hay (6 kg) and cubes (4 kg); Group 2: lucerne hay (6 kg) and cubes (4 kg); Group 3: tef hay (4 kg), cubes (6 kg) and fishmeal (200 g); Group 4: lucerne hay (6 kg), cubes (4 kg) and fishmeal (200 g). These 4 feeding regimens were chosen to obtain a wide range of dietary amino-acid contents. Tef hay is generally used as a source of roughage in large parts of South Africa. Its protein content is low compared to lucerne hay³⁴. The crude protein and amino-acid intake of mares in the different dietary groups has been described³⁵. The cubes were offered in 2 equal portions in the morning and afternoon³⁵. After the mares had foaled, the cubes were increased to 6 kg fed 3 times a day in equal portions and the hay to 7 kg, while the fishmeal remained at 200 g/day. Each mare and foal received their cubes separately, but during the morning feed, when the mares received their fishmeal, the foals were prevented from eating with the mares. Throughout the trial the mares finished all feed offered to them every day. At 3 months of age the foals received 1 kg of cubes and *c.* 2 kg of roughage per day. Each group of mares was kept in a 30 × 50 m paddock. The foaling date was recorded in all cases.

Daily teasing commenced 2 d after parturition and continued until 90 d of pregnancy. The ovaries of the mares in oestrus were palpated daily. The ovaries of mares that had not shown signs of oestrus by 10 d *post partum* were palpated weekly until signs of oestrus became evident. During examinations the size and consistency of the ovaries and follicles present were determined as described previously³⁶. A Squibb Model 2000 scanner with a 5 MHz probe was

^aDepartment of Human and Animal Physiology, University of Stellenbosch, Private Bag X1, Matieland, 7602 South Africa.

Received: December 1997. Accepted: June 1998.

Table 1: Comparison of the mean duration (days) from parturition to 1st oestrus and ovulation and the following oestrous period and follicular size (mm) during both ovulations after parturition in mares in 4 dietary groups.

Oestrous cycle	Dietary groups				Mean total
	1	2	3	4	
1st oestrus					
Number of mares	6	4 (5)	4	5 (6)	19
Days from parturition to 1st oestrus	7.0 ± 3.3	8.3 ± 1.5 (26.2 ± 35.8) ^a	9.7 ± 2.9	10.0 ± 3.5 (19.0 ± 20.3) ^a	8.7 ± 3.1
Days in oestrus	8.8 ± 3.7	5.0 ± 2.4	4.3 ± 0.9	5.3 ± 1.9	6.1 ± 3.0
Days from onset of oestrus to ovulation	7.3 ± 3.4	3.5 ± 1.7	3.3 ± 0.9	3.8 ± 2.3	4.7 ± 2.8
Days from parturition to 1st ovulation	14.1 ± 3.5	10.7 ± 3.0 (29.2 ± 37.2) ^a	12.8 ± 4.1	14.0 ± 5.6 (22.5 ± 21.4) ^a	13.2 ± 4.1
<i>Size of follicle (mm)</i>					
Onset of oestrus	23 ± 7.5	27 ± 2.9	23 ± 6.3	24 ± 5.8	24 ± 5.8
Day before ovulation	35 ± 7.3	36 ± 9.5	36 ± 7.5	37 ± 5.2	36 ± 6.7
2nd oestrus					
Number of mares	5	3	4	4	16
Days in oestrus	5.8 ± 3.7	3.7 ± 1.5	6.2 ± 1.7	6.8 ± 1.7	5.7 ± 2.5
Days from onset of oestrus to ovulation	5.0 ± 3.7	3.7 ± 1.5	5.2 ± 1.2	5.5 ± 2.4	4.9 ± 2.4
<i>Size of follicle (mm)</i>					
Onset of oestrus	28 ± 6.0	22 ± 10.0	30 ± 5.0	26 ± 11.0	28 ± 6.0
Day before ovulation	36 ± 4.0	35 ± 5.0	41 ± 3.0	42 ± 3.0	39 ± 5.0
Number of oestrous cycles	7	9	3	4	
Mean duration of oestrous cycles (days)	20.7	19.2	24.0	23.0	20.8
Range (days)	16–31	13–22	22–28	21–26	13–31

^aMares that did not show oestrus within 20 days of parturition.

used for all ultrasonographic examinations.

Mares were mated during the 1st oestrus following foal heat. If a mare did not show signs of oestrus within 12 d *post partum* she was mated during the 1st oestrous period. Mares were mated when a 30–50 mm, soft, thin-walled follicle was palpable and daily thereafter until ovulation occurred, after which mating ceased.

Blood samples were collected on the day of parturition and every 2nd day thereafter until signs of oestrus were evident. Samples were collected daily during oestrus, and then every 3rd day until the following oestrous period. This procedure was followed until pregnancy was diagnosed. Blood samples were collected and the serum stored as described previously³⁷.

All mares and foals were weighed on the day of foaling and weekly thereafter until the foals were weaned at 6 months.

Hormone assays

Total unconjugated serum progestagen

Serum samples were analysed for total unconjugated progestagens. The radio-immunoassay procedure was principally as described by Youssefnejadian *et al.*⁴⁰ using an antibody generated in sheep according to the method of Odell

*et al.*²⁴ against 11 α -hydroxyprogesterone hemi-succinate BSA¹² as modified by Faure¹⁰. In addition, the antibody shows significant cross-reactivity with α -pregnane-3,20 dione, and to a much lesser extent with other pregnane derivatives of progesterone that are known to occur in the pregnant mare¹⁸. Cross-reactivity with major adrenocortical C18 and C19 steroids is for all practical purposes non-existent³³.

Follicle-stimulating hormone (FSH)

FSH concentrations were determined using the Amerlex-M FSH (Code DM 3070/3071) kit (Amersham, UK). Use of this antibody was based on the results of Alexander *et al.*².

Luteinising hormone (LH)

LH concentrations were determined using the method described by Niswender *et al.*²³ as modified by Visser³⁹.

Statistical analysis

Analysis of variance was conducted using the LSML 76 computer programme¹⁵.

RESULTS

The mean body weight of mares on the day of foaling was: Group 1, 437 kg; Group 2, 472 kg; Group 3, 441 kg; Group 4, 454 kg. The mean birth weight of the foals of the mares in each group was:

Group 1, 47.3 kg; Group 2, 47.2 kg; Group 3, 51.2 kg; Group 4, 46.2 kg.

The number of days from parturition until 1st signs of oestrus (foal heat), the duration of the oestrus periods, time of ovulation, diameter of follicles and the duration of the oestrous cycle following foal heat are shown in Table 1 for the lactating mares in the 4 dietary groups. The results that include the 2 mares (No. 8 in Group 2 and No. 20 in Group 4) that did not show signs of foal heat or oestrus until 80 d and 64 d *post partum* respectively, are shown separately in Table 1.

No differences in the diameter (mm) of the largest follicle at the onset of oestrus and the diameter of the largest follicle on the day before ovulation were found between groups.

In Table 2 a comparison is made between the onset of 1st signs of oestrus and the time of ovulation in mares that foaled during August and September (transitional period) and mares that foaled during October to December (physiological breeding season), irrespective of their nutritional group. Those that foaled in the transitional period showed the 1st signs of oestrus on average 24 d *post partum*, compared to 9 d in those that foaled in the physiological breeding season, and on average the 1st ovulations occurred 27.1 d and 14.6 d *post partum*

Table 2: Comparison of the mean number of days from parturition until the first signs of oestrus in mares that foaled early in the season (August - September) and mares that foaled later in the season (October - December).

Date of foaling	Number of mares	Days from foaling to 1st oestrus	Days from foaling to 1st ovulation	Duration of 1st oestrous period	Diameter of follicle (mm)	
					At onset	At ovulation
14 Aug – 28 Sept	8	24.0 ± 30.0 (5–80)	27.1 ± 30.2 (6–64)	5.1 ± 2.6 (2–9)	25 ± 5.0 (15–30)	38 ± 7.0 (30–50)
4 Oct – 15 Dec	12	9.0 ± 3.5 (4–14)	14.6 ± 3.2 (8–19)	6.8 ± 3.2 (4–14)	24 ± 6.0 (10–30)	35 ± 6.0 (25–40)




Fig. 1: Four mean serum progesterone concentration (ng/ml) patterns identified from parturition to ovulation at foal heat: Days 6, 9–11, 12–13 and 15–19.

respectively. The mean duration of the 1st oestrous period for the early and late foalers was 5.1 d and 6.8 d respectively.

No differences in serum progesterone concentrations were found during the time from parturition to foal heat among the 4 nutritional groups. The mean serum progesterone concentrations of mares ($n = 18$) that ovulated on Days 6, 9–11, 12–13 and 15–19 *post partum* are shown in Fig. 1.

The progesterone concentrations of mares during the 1st oestrous cycle after foal heat did not differ between the nutritional groups, and therefore the results of mares in all 4 dietary groups were pooled. During foal heat, serum progesterone concentrations increased within 2 days post-ovulation. Serum progesterone concentrations were 7–20 ng/ml at 4–6 days post-ovulation and were maintained at these concentrations for 13–15 days, after which they decreased to reach baseline values *c.* 2 days before the next ovulation. After pooling the results, 4 different

patterns of progesterone secretion were identified according to the duration of the interovulatory period, namely A: 17–19 d; B: 20–22 d; C: 24–25 d, and D: 29–30 d (Fig. 2).

Persistent CLs were identified in 2 mares. Mare No. 8 (Group 2: lucerne hay and cubes), which foaled in mid-August, ovulated on Day 9 *post partum* without showing signs of oestrus for the following 80 days and had low (<5 ng/ml) serum progesterone concentrations during this period (Fig. 3). Mare No. 20 (Group 4: lucerne hay, cubes and fishmeal), which foaled in September, showed no signs of oestrus but ovulated on Day 11 *post partum*. She then maintained low (<5 ng/ml) serum progesterone concentrations until signs of oestrus and she ovulated on Day 66 *post partum* (Fig. 3).

Serum LH and FSH concentrations were established only in mares in Group 1 (lowest quality protein) and Group 4 (highest quality protein), from parturition

until foal heat and the following oestrous cycle until after ovulation. No differences in the LH concentrations between dietary groups that could be attributed to protein nutrition were found, and therefore the results were pooled to demonstrate the general secretion pattern. The mean pooled LH and progesterone concentrations from parturition until after the 2nd ovulation are shown in Fig. 4. The mean duration from foaling to ovulation at foal heat was 13.7 d. LH concentrations increased gradually from Day 6 *post partum* until 2 days before ovulation, when the LH concentrations began to peak. In 11 of the 18 mares the LH peak occurred on the day of ovulation, and in the remaining 8 during the following day. LH concentrations decreased to baseline levels within 48 h post-ovulation.

The serum FSH concentrations were pooled, as no differences were found between mares in the 4 dietary groups from parturition to 1st ovulation (Fig. 5).

Table 3: Conception rates of mares in 4 dietary groups.

Dietary group	Oestrous period after parturition when mares were mated						Mated oestrous period per conception
	1st ^a	2nd	3rd	4th	5th	6th	
1	1 (100 %) ^b	5 (80 %)	1 (0 %)	1 (0 %)	—	—	1.6
2	—	3 (33 %)	3 (33 %)	2 (50 %)	1 (0 %)	1 (100 %)	2.5
3	—	4 (100 %)	—	—	—	—	1.0
4	—	5 (100 %)	—	—	—	—	1.0

^aFoal heat.

^bNumber of mares (conception rate of mares mated during cycle).




Fig. 2: Mean serum progesterone concentration (ng/ml) in mares with interovulatory periods of (A) 17–19 d (n = 4), (B) 20–22 d (n = 5), (C) 24–25 d (n = 5) and (D) 29–30 d (n = 2) during the oestrous cycle following foal heat.




Fig. 3: Serum progesterone concentrations (ng/ml) of Mare 8 (Group 1) and Mare 20 (Group 4) that developed persistent corpora lutea after ovulation.




Fig. 4: Mean serum progesterone concentration (ng/ml) of mares (n = 18) that ovulated at foal heat followed by a normal oestrous cycle.




Fig. 5: Mean serum FSH concentration (ng/ml) of mares (n = 18) that ovulated at foal heat followed by a normal oestrous cycle.

At parturition the mean FSH concentrations were relatively high (25 ng/ml) and increased (30 ng/ml) during the 1st 3 days *post partum*. In all mares that showed foal heat, FSH concentrations declined after Day 3 to reach minimum concentrations (15 ng/ml) by Day 8. The mean FSH concentrations increased at the onset of foal heat and reached high concentrations (>30 ng/ml) the time of ovulation. Thereafter concentrations declined 1–2 d after

ovulation to reach low values (<20 ng/ml) 5–7 d after ovulation. An increase in FSH concentrations with maximum concentrations of approximately 33 ng/ml were found from Days 9–12 of the following cycle, after which the concentrations declined to minimum values (<20 ng/ml) by Day 18. Increases were recorded 2–3 d before the next ovulation (Fig. 5).

The conception rate of the lactating mares is shown in Table 3. Mares in

Groups 3 and 4 conceived at the 1st mated oestrus period compared to the 1.6 (Group 1) and 2.5 (Group 2) mated oestrous periods per conception in the groups receiving the low-quality protein diets.

DISCUSSION

The total daily protein requirement of a mare with a body weight of 500 kg is 800–870 g/day during late pregnancy and

1284–1427 g during the 1st 3 months of lactation²⁰. In the current study the mean body weight of all the mares in the respective dietary groups was below 500 kg immediately after foaling. The total daily crude protein intake of the mares in the 4 dietary groups varied between 1000 and 1800 g/day during late pregnancy, while the lowest daily crude protein intake of 1200 g received by the mares in Group 1 during lactation is considered to meet their requirements³⁵. It is thus clear that, according to the recommendations of the National Research Council, Washington DC²⁰, none of the mares suffered from a protein deficiency during any stage of this trial.

Foal heat normally commences 7–12 d *post partum*^{14,22}. Although there was no significant difference in the number of days from parturition until the onset of foal heat between groups, the mean duration of this period for mares in Group 1 (low-quality protein) was 7 d compared to 10 d in Group 4 (high-quality protein). The 2 mares that showed the 1st signs of oestrus 80 and 64 days after foaling, foaled early during the season (transitional phase) and the prolonged *post partum* anoestrous periods were probably due to the effect of season/photoperiod and not protein nutrition. The effect of season on the incidence of foal heat has been described in Thoroughbred and Percheron mares³⁰. A higher incidence of failure to show foal heat was found amongst mares that foaled during early spring and also amongst high milk producers³⁰. The pooled results (Table 2) indicated that the time of foaling (season) played an important role in the incidence of oestrus. In a previous study with barren mares kept in the same locality, only 10 % of the mares ovulated between 29 August and 20 September, while 70 % ovulated from the last week in September to mid-October³⁶. When the 2 mares described above are excluded, the current results are in agreement with the literature^{14,21,36}. Protein nutrition apparently had very little effect on the duration of the period from parturition to the onset of foal heat in the mares used in this study.

The average duration of foal heat (Table 1) of 8.8 d for Group 1 (low-quality protein) was approximately 3–4 d longer than that recorded in any of the other groups. Mares in Group 1 ovulated on average 7.3 d after the onset of oestrus, approximately 3 d later than the mares in the other groups. However, when the number of days from parturition to ovulation is compared, no differences were found among groups (Table 1).

Table 1 shows that the duration of

oestrous cycles following foal heat was 3–4 d longer in the fishmeal-supplemented Groups (3 and 4) than in Groups 1 and 2, which did not receive fishmeal.

In addition, the duration of the oestrous cycles in the mares in Groups 3 and 4 showed less variation (22–28 d) than mares in Groups 1 and 2, where a range of 13–31 days was recorded. This might indicate improved endocrine function with regard to the production of progesterone and possibly LH, which is important for early CL stimulation after ovulation. This is in agreement with previous results, where high-quality protein decreased the time to 1st ovulation after the anovulatory period in barren mares^{36,37}. However, the mean duration of the oestrous cycles was 20.8 d (13–31), which compares favourably with the 20.7 d reported for Thoroughbred mares in the southern hemisphere³⁰.

During late pregnancy and at parturition the brain-pituitary-ovarium (BPO) axis and the uterus are exposed to high circulating concentrations of progesteragen, oestrogen, prostaglandins and oxytocin. After parturition, not only must the uterus recover, but also the BPO axis of the positive and negative endocrine feedback systems to enable resumption of the ovarian cycle³². Total serum progesteragen concentrations did not differ between groups from parturition until the onset of foal heat. The relatively high total serum progesteragen concentrations (5.2–16.4 ng/ml) found on the day of parturition declined rapidly to concentrations of 2–3 ng/ml within the next 48 h, which is in agreement with previous findings^{13,14}. Although the effect of protein nutrition on progesteragen concentration during the *post partum* period is uncertain, it appears that in the mares that only came into oestrus 15–19 d *post partum* the progesteragen concentrations remained at between 1.5–3.0 ng/ml until 12–14 days *post partum*. The possible suppression of GnRH secretion by this persistently elevated level of serum progesteragen is unknown at this stage.

The serum progesteragen secretion patterns as described and shown in Fig. 2 during the oestrous cycle following foal heat are in agreement with those described previously^{3,31}. The mean serum progesteragen concentrations of mares that exhibited interovulatory periods of 17–19 d, 20–22 d, 24–25 d and 29–30 d (Fig. 2) indicate that, as the duration of the oestrous cycle increases, mid-cycle progesteragen concentrations remain elevated for a longer period during the luteal phase of the cycle, and that maximum progesteragen concentrations occur 2–3 d later

during the oestrous cycle (Fig. 2). This might initially be the result of under-stimulation of the luteal tissue by LH, leading to lower production of progesterone. Early embryonic loss before Day 12 has no prolonged effect on the duration of the oestrous cycle¹¹.

The 2 mares that ovulated 6–11 days *post partum* and developed persistent *corpora lutea* only came into oestrus between 40–80 days after the 1st ovulation. Both mares foaled during the transitional period (August–September), at the time of the year when the incidence of ovulation amongst barren mares is reported to be low³⁶. A similar decline in ovarian activity after parturition as well as ovulation during foal heat in mares that foaled during winter or early spring has been described²⁶. However, the results of this study indicate that the protein intake and quality had very little effect on progesterone production during the early *post partum* period.

No differences in the time of the LH peak secretion or the maximum concentration were found between dietary Groups 1 and 4 during ovulation at foal heat or during the following oestrous period. It is accepted that the photoperiod might have influenced LH secretion, as has been described³⁷. Current results are in agreement with previous reports of low LH concentrations at parturition²⁶. LH concentrations are reported to increase gradually from Day 6 *post partum* until 2 days before ovulation, followed by a sharp increase to reach peak concentrations just after ovulation²⁶. As shown in Fig. 4, the increase in LH coincides with low progesteragen concentrations at 4–6 days *post partum*. Serum oestrogen concentrations that increase 5–10 days *post partum*²¹ stimulate secretion of LH. According to the results of the individual mares in this study, the LH peak was recorded on the day of ovulation or the day following ovulation, which is in agreement with current literature²⁶.

Although no differences in the concentrations of FSH were found among mares in the different dietary groups, the observed decrease in FSH concentrations after parturition and the subsequent increase before oestrus and ovulation during foal heat have been described previously^{14,26}. It is evident from the results of the individual mares that a relationship exists between the time when FSH concentration begins to decline and the time of ovulation. It therefore indicates that in cases where foal heat is delayed, the typical FSH secretion pattern does not occur at the same early stage. A similar stimulatory effect of high-quality

dietary protein on FSH production found in barren mares³⁷ was not observed in this group of mares during the *post partum* period. The overriding effect of season/photoperiod over protein quality of the diet might have been the reason for this finding.

The 10–14-day cyclical secretion pattern of FSH found in the lactating mares in this study (Fig. 5) is in agreement with previous reports^{9,27,37}. FSH stimulates follicular growth and a FSH peak normally occurs a few days after ovulation. Oestrogen produced by the developing follicles stimulates the LH surge and subsequently ovulation. In Equidae the mid-cycle FSH peak results in follicular development and the formation of mid-cycle follicles. These follicles generally undergo regression as LH concentrations, which increase slightly during this period but do not reach peak concentrations, are insufficient to stimulate follicular maturation and ovulation²⁹.

Only 56 % of the mares that were not supplemented with fishmeal (Groups 1 and 2) conceived when mated during the oestrous period following foal heat, compared to the 100 % conception rate in the mares in Groups 3 and 4 (fishmeal supplemented). However, all mares mated in Groups 1 and 2 eventually conceived during subsequent oestrous periods.

CONCLUSION

Inclusion of high-quality protein in the diet of lactating mares resulted in an increased conception rate. The incidence of abnormally long or short oestrous cycles was reduced in mares receiving fishmeal supplementation. Serum progesterone, FSH and LH concentrations were unaffected by protein intake and quality. It would appear that the normal endocrine function of the brain-pituitary-ovarian axis is resumed within 6–18 days *post partum* with the possible exception of mares foaling during early spring. The effect of season/photoperiod influenced the *post partum* breeding results of mares that foaled during early spring detrimentally. Uterine involution may be delayed in mares fed low-quality protein diets resulting in a decrease in conception rate, an increase in the number of services per conception and, more importantly, in Thoroughbreds an increased possibility of lowered pregnancy rates at the end of the restricted breeding season.

ACKNOWLEDGEMENTS

We thank Mr S W P Cloete, Elsenburg Agricultural Development Institute, for statistical analysis and the S A Medical Service for providing experimental animals and financial support.

REFERENCES

1. Achnelt E, Plas J 1947 Time of the first ovulation in mares after foaling. *Animal Breeding Abstracts* 15: 233
2. Alexander S L, Irvine C H G, Turner J E 1987 Comparison by three different radio-immuno assay systems of the polymorphism of plasma FSH in mares in various reproductive states. *Journal of Reproduction and Fertility Supplement* No. 35: 9–18
3. Allen W R 1987 Endocrinological and immunological aspects of pregnancy and pregnancy loss. *Proceedings, 9th Bain Fallon Lectures Sydney*: 61–74
4. Bain A M 1957 Estrus and infertility of the Thoroughbred mare in Australasia. *Journal of the American Veterinary Medical Association* 131: 179–185
5. Belonje P C, Van Niekerk C H 1975 A review of the influence of nutrition upon the oestrous cycle and early pregnancy in the mare. *Journal of Reproduction and Fertility Supplement* No. 23: 167–169
6. Deskur S 1965 The first oestrus and ovulation after parturition in the mare. *Animal Breeding Abstracts* 33: 533
7. Doreau M, Martin-Rosset W, Boulot S 1988 Energy requirements and the feeding of mares during lactation: review. *Livestock Production Science* 20: 53–68
8. Du Plessis J L 1964 Some observations and data in Thoroughbred breeding. *Journal of the South African Veterinary Medical Association* 35: 215–221
9. Evans M J, Irvine C H G 1975 Serum concentrations of FSH, LH and progesterone during the oestrous cycle and early pregnancy in the mare. *Journal of Reproduction and Fertility Supplement* No. 23: 193–200
10. Faure A S 1975 Vroeë embrionale verliese weens wanvoeding by Merinoskape. MSc thesis, University of Stellenbosch
11. Forde D, Keenan L, Wade J, O'Connor M, Roche J F 1987 Reproductive wastage in the mare and its relationship to progesterone in early pregnancy. *Journal of Reproduction and Fertility Supplement* No. 35: 493–495
12. Furr B J 1973 Radio-immunoassay of progesterone in peripheral plasma of the domestic fowl venous plasma. *Acta Endocrinologica* 72: 89–100
13. Ganjam V K, Kenny R M, Flickinger G 1975 Effect of exogenous progesterone on its endogenous levels: biological half-life of progesterone and lack of progesterone binding in mares. *Journal of Reproduction and Fertility Supplement* No. 23: 183–188
14. Ginther O J 1979 *Reproductive biology of the mare. Basic and applied aspects*. McNaughton and Gunn, Ann Arbor, Michigan
15. Harvey W R 1977 *Users guide for LSML 76 mixed model least squares and maximum likelihood computer program*. Ohio State University, Ohio
16. Henneke D R, Kreider J L 1979 Effects of restrictive suckling on post partum reproductive performance in mares. *American Journal of Veterinary Research* 40: 1281–1284
17. Henneke D R, Potter G D, Kreider J L 1981 Breeding efficiency in mares fed different levels of energy during late gestation. *Proceedings, 7th Equine Nutrition Physiology Symposium*. Virginia State University, Petersburg: 101–104
18. Holtan D W, Nett T M, Estergreen V L 1975 Plasma progestins in pregnant, post partum and cycling mares. *Journal of Animal Science* 40: 251–260
19. Matthews R G, Ropiha R T, Butterfield R M 1967 The phenomenon of foal heat in mares. *Australian Veterinary Journal* 43: 579–582
20. National Research Council 1989 *Nutrient requirements of horses* (5th revised edn). National Academy Press, Washington DC
21. Nett T M, Holtan D W, Estergreen V L 1975 Levels of LH, prolactin and oestrogens in the serum of post-partum mares. *Journal of Reproduction and Fertility Supplement* No. 23: 201–206
22. Nishikawa Y 1959 *Studies on reproduction in horses*. Japan Racing Association Minotoku, Tokyo
23. Niswender H D, Reichert L E, Midgley A R, Nalbandov A V 1969 Radio-immunoassay for bovine and ovine luteinizing hormone. *Endocrinology* 84: 1166
24. Odell W D, Abraham G A, Skowsky W R, Hescocox M A, Fisher D A 1971 Production of antisera for radio-immunoassays. In Odell W D, Daughaday W H (eds) *Principles of competitive protein binding assays*. J B Lippincott & Co, Philadelphia: 57–76
25. Pagan J D, Hintz H F, Rounsaville T R 1984 The digestible energy requirements of lactating pony mares. *Journal of Animal Science* 58: 1382–1387
26. Palmer E, Draincourt M A 1983 Some interactions of season of foaling, photoperiod and ovarian activity in the equine. *Livestock Production Science* 10: 197–210
27. Pineda M H, Ginther O J 1972 Inhibition of estrus and ovulation in mares treated with an antiserum against an equine pituitary fraction. *American Journal of Veterinary Research* 33: 1775–1780
28. Tolkdorff E, Jochle W, Lamond D R, Klug E, Merkt H 1976 Induction of ovulation during the post-partum period in the Thoroughbred mare with a prostaglandin analogue synchrocept. *Theriogenology* 6: 403–412
29. Vandeplasse M, Henry M, Coryn M 1979 The mature mid-cycle follicle in the mare. *Journal of Reproduction and Fertility Supplement* No. 27: 157–162
30. Van Niekerk C H 1965 The breeding cycle, ovarian changes and tubal sojourn of ova in the mare. M Med Vet thesis, University of Pretoria
31. Van Niekerk C H 1973 Morphology and physiological changes in the genital system of mares. DVSc thesis, University of Pretoria
32. Van Niekerk C H 1976 Limitations to female reproductive efficiency. In G J Tomes, D E Robertson, R J Lightfoot (eds) *Proceedings of the International Congress on Sheep Breeding, Perth, Australia, August 1976*: 294–309
33. Van Niekerk C H, Morgenthal J C 1982 Fetal loss and the effect of stress on plasma progesterone levels in pregnant Thoroughbred mares. *Journal of Reproduction and Fertility Supplement* No. 32: 453–457
34. Van Niekerk F E, Van Niekerk C H 1997 The effect of dietary protein on reproduction in the mare. I. The composition and evaluation of the digestibility of dietary protein from different sources. *Journal of the South African Veterinary Association* 68: 78–80
35. Van Niekerk F E, Van Niekerk C H 1997 The effect of dietary protein on reproduction in the mare. II. Growth of foals, body mass of mares and serum protein concentrations of mares during the anovulatory, transitional and pregnant periods. *Journal of the South African Veterinary Association* 68: 81–85
36. Van Niekerk F E, Van Niekerk C H 1997 The effect of dietary protein on reproduction in the mare. III. Ovarian and uterine changes

during the anovulatory, transitional and ovulatory periods in the non-pregnant mare. *Journal of the South African Veterinary Association* 68: 86–92

37. Van Niekerk F E, Van Niekerk C H 1997 The effect of dietary protein on reproduction in the mare. IV. The serum progestagen, FSH, LH and melatonin concentrations of non-

pregnant mares during the anovulatory and transitional periods. *Journal of the South African Veterinary Association* 68: 114–120

38. Van Niekerk C H, Van Heerden J S 1972 Nutrition and ovarian activity of mares early in the breeding season. *Journal of the South African Veterinary Association* 43: 355–360

39. Visser J H 1991 'n Geslagsendokrinologiese ondersoek na die seisoenaliteit van ramme van vyf verskillende skaap- en bokrasse. MSc thesis, University of Stellenbosch

40. Yousefnajadian E, Florensa E, Collins W P, Sommerville I F 1972 Radio-immunoassay of plasma progesterone. *Journal of Steroid Biochemistry* 3: 893